



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Reply to 'Misestimation of heritability and prediction accuracy of male-pattern baldness'

Citation for published version:

Pirastu, N, Joshi, PK, de Vries, PS, Cornelis, MC, Keum, N, Franceschini, N, Colombo, M, Giovannucci, EL, Spiliopoulou, A, Franke, L, North, KE, Kraft, P, Morrison, AC, Esko, T & Wilson, JF 2018, 'Reply to 'Misestimation of heritability and prediction accuracy of male-pattern baldness'', *Nature Communications*, vol. 9, no. 1, pp. 2538. <https://doi.org/10.1038/s41467-018-04808-2>

Digital Object Identifier (DOI):

[10.1038/s41467-018-04808-2](https://doi.org/10.1038/s41467-018-04808-2)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Publisher's PDF, also known as Version of record

Published In:

Nature Communications

Publisher Rights Statement:

This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



CORRESPONDENCE

DOI: 10.1038/s41467-018-04808-2

OPEN

Reply to 'Misestimation of heritability and prediction accuracy of male-pattern baldness'

Nicola Pirastu¹, Peter K. Joshi¹, Paul S. de Vries², Marilyn C. Cornelis³, NaNa Keum⁴, Nora Franceschini⁵, Marco Colombo⁶, Edward L. Giovannucci^{4,7,8}, Athina Spiliopoulou^{6,9}, Lude Franke¹⁰, Kari E. North⁵, Peter Kraft¹¹, Alanna C. Morrison², Tõnu Esko^{12,13} & James F. Wilson^{1,14}

Yap et al.¹ present two criticisms of our recent analysis² of the genetic architecture of male pattern baldness (MPB)². First they note our earlier study in ref.² overestimated SNP heritability (hereafter heritability) by excluding people in category two on the UK Biobank scale. We agree, heritability should have been reported as 0.64 not 0.94. This arose for a natural, if mistaken for this purpose, desire to categorize subjects clearly on a binary scale, and thus exclude indeterminate subjects. Their second criticism is that we overestimated the proportion of heritability explained by the 71 loci that have been identified. We stand by the broad conclusion that about one-third of the genetic effects (on a baldness trait dichotomized as category 1 versus 2, 3, or 4) are explained by the 71-locus SNP score.

In principle, the proportion of polygenic variance explained by the SNP score can be evaluated in the following three possible ways: (1) as the ratio of the phenotypic variance explained by the SNP score to the variance explained by polygenic effects; (2) as the ratio between the heritability due to the SNPs and the baseline heritability estimate; or (3) as ratio of the reduction in polygenic variance in a model that includes the SNP score to the polygenic variance in a model that does not include this fixed effect (see Supplementary Method for details). These three methods should give the same result if the residual variance and the phenotypic variance do not change between the models with and without the SNPs.

Yap et al. have used the first method, and estimate that the 107 SNPs from 71 loci explain about 15–20% of variation in total liability. Our own estimate using the same method is 20% on the

liability scale, close to theirs, implying that about 31% of the total heritability of 0.61 is explained by the SNP score. Our article, however, reported an estimate by method (2), in which the ratio of the difference in heritability in models including and excluding the SNP to the baseline heritability was 38%. Including category two did not change this estimate (Supplementary Table 1 method (2)). Of course, to evaluate predictive performance requires an independent test dataset, beyond the scope of both our original study² and the correspondence¹.

GCTA implements a mixed linear model and therefore estimates phenotypic variance from the variances of the random effects in the model. Therefore, the estimated phenotypic variances from models with different fixed effects (i.e., with and without the SNP predictor) are different. We thus applied method (3) which is not affected by the same issue as it works on the absolute and not relative scale (see Supplementary Method) and gives an even greater estimate of 45%. Given the limitations of fitting mixed linear models to a binary trait to estimate the parameter of interest, it is not easy to be certain which is the best one. However, irrespective of which is used, our conclusion that we can explain a relatively large proportion of heritability using SNPs from only 71 loci is still valid.

Having now corrected the error in the estimation of heritability, we thus believe that the remainder of the results and conclusions are still valid, including in particular that we can explain a large proportion of the genetic variance using a relatively small number of SNPs. Furthermore, our identification and replication of several new loci for MBP remains accurate and

¹ Centre for Global Health Research, Usher Institute of Population Health Sciences and Informatics, University of Edinburgh, Teviot Place, Edinburgh EH8 9AG, UK. ² Human Genetics Center, University of Texas Health Science Center at Houston, Houston, TX 77030, USA. ³ Department of Preventive Medicine, Northwestern University Feinberg School of Medicine, Chicago, IL 60611, USA. ⁴ Department of Nutrition, Harvard T. H. Chan School of Public Health, Boston, MA 02115, USA. ⁵ Department of Epidemiology and Carolina Center for Genome Sciences, University of North Carolina, Chapel Hill, NC 27599, USA. ⁶ Center for Population Health Sciences, Usher Institute of Population Health Sciences and Informatics, University of Edinburgh, Teviot Place, Edinburgh EH8 9AG, UK. ⁷ Department of Epidemiology, Harvard T. H. Chan School of Public Health, Boston, MA 02115, USA. ⁸ Channing Division of Network Medicine, Department of Medicine, Brigham and Women's Hospital and Harvard Medical School, Boston, MA 02115, USA. ⁹ Pharmatics Ltd, Edinburgh EH16 4UX, UK. ¹⁰ Department of Genetics, University Medical Center, 9713 GZ Groningen, The Netherlands. ¹¹ Program in Genetic Epidemiology and Statistical Genetics, Harvard T. H. Chan School of Public Health, Boston, MA 02115, USA. ¹² Estonian Genome Center, University of Tartu, 51010 Tartu, Estonia. ¹³ Broad Institute of Harvard and MIT, Cambridge, MA 02142, USA. ¹⁴ MRC Human Genetics Unit, Institute of Genetics and Molecular Medicine, University of Edinburgh, Western General Hospital, Crewe Road, Edinburgh EH4 2XU, UK. Correspondence and requests for materials should be addressed to N.P. (email: nicola.pirastu@ed.ac.uk)

increases the understanding and biological interpretation whilst highlighting the shared genetics with other traits.

Data availability

All data that support the findings of this study are available from the corresponding author upon reasonable request.

Received: 8 March 2018 Accepted: 12 April 2018

Published online: 29 June 2018

References

1. Yap, C. X., et al. Misestimation of heritability and prediction accuracy of male-pattern baldness. *Nat. Commun.* <https://doi.org/10.1038/s41467-018-04807-3> (2018).
2. Pirastu, N., Joshi, P. K. & de Vries, P. S. et al. GWAS for male-pattern baldness identifies 71 susceptibility loci explaining 38% of the risk. *Nat. Commun.* **8**, 1584 (2017).

Author contributions

N.P., J.F.W., and P.J. wrote the manuscript. N.P. performed the statistical analyses. All authors critically reviewed the manuscript.

Additional information

Supplementary Information accompanies this paper at <https://doi.org/10.1038/s41467-018-04808-2>.

Competing interests: The authors declare no competing interests.

Reprints and permission information is available online at <http://npg.nature.com/reprintsandpermissions/>

Publisher's note: Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2018